


Convenient B.I.D. Gantanol[®] (sulfamethoxazole)



Before prescribing, please consult complete product information, a summary of which follows:

Indications: Acute, recurrent or chronic nonobstructed urinary tract infections (primarily pyelonephritis, pyelitis and cystitis) due to susceptible organisms. **Note: Carefully coordinate *in vitro* sulfonamide sensitivity tests with bacteriologic and clinical response; add aminobenzoic acid to follow-up culture media. The increasing frequency of resistant organisms limits the usefulness of antibacterials including sulfonamides, especially in chronic or recurrent urinary tract infections.** Measure sulfonamide blood levels as variations may occur; 20 mg/100 ml should be maximum total level.

Contraindications: Sulfonamide hypersensitivity; pregnancy at term and during nursing period; infants less than two months of age.

Warnings: Safety during pregnancy has not been established. Sulfonamides should not be used for group A beta-hemolytic streptococcal infections and will not eradicate or prevent sequelae (rheumatic fever, glomerulonephritis) of such infections. Deaths from hypersensitivity reactions, agranulocytosis, aplastic anemia and other blood dyscrasias have been reported and early clinical signs (sore throat, fever, pallor, purpura or jaundice) may indicate serious blood disorders. Frequent CBC and urinalysis with microscopic examination are recommended during sulfonamide therapy. Insufficient data on children under six with chronic renal disease.

Precautions: Use cautiously in patients with impaired renal or hepatic function, severe allergy, bronchial asthma; in glucose-6-phosphate dehydrogenase-deficient individuals in whom dose-related hemolysis may occur. Maintain adequate fluid intake to prevent crystalluria and stone formation.

Adverse Reactions: *Blood dyscrasias* (agranulocytosis, aplastic anemia, thrombocytopenia, leukopenia, hemolytic anemia, purpura, hypoprothrombinemia and methemoglobinemia); *allergic reactions*

(erythema multiforme, skin eruptions, epidermal necrolysis, urticaria, serum sickness, pruritus, exfoliative dermatitis, anaphylactoid reactions, periorbital edema, conjunctival and scleral injection, photosensitization, arthralgia and allergic myocarditis); *gastrointestinal reactions* (nausea, emesis, abdominal pains, hepatitis, diarrhea, anorexia, pancreatitis and stomatitis); *CNS reactions* (headache, peripheral neuritis, mental depression, convulsions, ataxia, hallucinations, tinnitus, vertigo and insomnia); *miscellaneous reactions* (drug fever, chills, toxic nephrosis with oliguria and anuria, periarteritis nodosa and L.E. phenomenon). Due to certain chemical similarities with some goitrogens, diuretics (acetazolamide, thiazides) and oral hypoglycemic agents, sulfonamides have caused rare instances of goiter production, diuresis and hypoglycemia as well as thyroid malignancies in rats following long-term administration. Cross-sensitivity with these agents may exist.

Dosage: *Systemic sulfonamides are contraindicated in infants under 2 months of age* (except adjunctively with pyrimethamine in congenital toxoplasmosis).

Usual adult dosage: 2 Gm (4 tabs or teasp.) initially, then 1 Gm *b.i.d.* or *t.i.d.* depending on severity of infection.

Usual child's dosage: 0.5 Gm (1 tab or teasp.)/20 lbs of body weight initially, then 0.25 Gm/20 lbs *b.i.d.* Maximum dose should not exceed 75 mg/kg/24 hrs.

Supplied: Tablets, 0.5 Gm sulfamethoxazole; Suspension, 0.5 Gm sulfamethoxazole/teaspoonful.



Roche Laboratories
Division of Hoffmann-La Roche Inc.
Nutley, New Jersey 07110